

XP-002543705

C:\EPOPROGS\SEA\1.1.1.\epodata\sea\eplog\internal.log

WPI / Thomson

AN - 1983-26776K [11]
 AP - JF197301211394 19731029
 CPY - MATS-I
 DC - B04
 DW - 198311
 IC - A61K35/78
 LNKA- 1983-026238
 M1 - [01] M423 M781 M903 P816 V400 V404
 MC - B04-A07D B12-H05
 PA - (MATS-I) MATSUI T
 PH - JF58009084B B 19830218 DW198311
 JF50070512 A 19750612 DW198311
 PR - JF197301211394 19731029
 XIC - A61K-035/78; A61K-036/185; A61K-036/60; A61P-003/00; A61P-003/08;
 C12C-003/00; C12C-007/00
 AB - Cones of hop obtd. within 35-45 days from flowering are dried
 naturally without exposure to sun or by exposing to air at 40-60 deg.C
 for 5-15 hrs., adjusting moisture content to 10-20%. The cones are
 crushed after drying, and the prod. used in the treatment of diabetes.
 (J50070512)
 ICAI- A61K36/60; A61P3/08; C12C3/00; C12C7/00
 ICCI- A61K36/185; A61P3/00; C12C3/00; C12C7/00
 IW - POWDER HOP EXTRACT PREPARATION TREAT DIABETES
 IWW - POWDER HOP EXTRACT PREPARATION TREAT DIABETES
 NC - 1
 NPH - 2
 OPD - 1973-10-29
 PAW - (MATS-I) MATSUI T
 PD - 1975-06-12
 TI - Powdered hop extract prepn. - used for treatment of diabetes (J5
 12.6.75)